

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

D3

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

International Patent Classification<sup>5</sup>:

C07C 233/00

A1

(11) International Publication Number:

WO 94/12466

(43) International Publication Date:

9 June 1994 (09.06.94)

(21) International Application Number: PCT/US93/11625

(22) International Filing Date: 30 November 1993 (30.11.93)

(30) Priority Data:

103,932

30 November 1992 (30.11.92) IL

(71) Applicant (for all designated States except US): YISSUM  
RESEARCH DEVELOPMENT COMPANY OF THE  
HEBREW UNIVERSITY OF JERUSALEM [IL/IL]; 46  
Jabotinsky Street, 91 042 Jerusalem (IL).(71)(72) Applicant and Inventor: DEVANE, William, A. [US/US];  
5480 Wisconsin Avenue, Apt. 1410, Chevy Chase, MD  
20815 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): MECHOULAM, Raphael  
[IL/IL]; 12 Tchernichovsky Street, 92 581 Jerusalem (IL).  
BEUER, Aviva [IL/IL]; 18/10 Neva Shaanan Street, 93 707  
Jerusalem (IL). HANUS, Lumir [IL/IL]; 26/6 Brazil Street,  
96 784 Jerusalem (IL).(74) Agents: FLEIT, Martin et al.; Keck, Mahin & Cate, P.O. Box  
06110, Chicago, IL 60606-0110 (US).(81) Designated States: AU, CA, CZ, HU, JP, PL, SK, US,  
European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR,  
IE, IT, LU, MC, NL, PT, SE).

Published

With international search report.

(54) Title: FATTY ACID DERIVATIVES AND PHARMACEUTICAL COMPOSITIONS CONTAINING THEM

(57) Abstract

Pure polyunsaturated fatty acid amides and their derivatives. These synthetically produced compounds are able to mimic naturally occurring anandamides in the brain and bind the cannabinoid receptor. The compounds exhibit physiological activity and are useful as active ingredients in pharmaceutical compositions for the treatment of inflammation, migraines, spasticity activity, glaucoma, multiple sclerosis. The active compounds can be provided in isotope-tagged form.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

FATTY ACID DERIVATIVES AND PHARMACEUTICAL COMPOSITIONS CONTAINING THEM  
FIELD OF THE INVENTION

The invention relates to certain polyunsaturated fatty acid amides, and derivatives of these. Part of these are  
5 present in the brain, and part are products of synthesis. The novel pure compounds have a variety of pharmacological properties. They inhibit the specific binding of a cannabinoid probe to synaptosomal membranes. The compounds of the invention can be provided in radioactivity tagged  
10 form.

BACKGROUND OF THE INVENTION

Arachidonic acid ethanolamide (anandamide) and similar compounds are constituents of the brain. Anandamide and certain of the compounds similar with same, bind to the  
15 cannabinoid receptor. The binding of the anandamide to the cannabinoid receptor is similar to the binding of  $\Delta^9$ -tetrahydrocannabinol. There exist in the body many mediators, which are derivatives of arachidonic acid, such as prostaglandins and leukotrienes, which are present as  
20 large families of related compounds. Certain of these do not bind to the cannabinoid receptor, and it was one of the aims of the present invention to provide and identify compounds which have pharmacological properties similar to the properties of anandamide.

25 The existence of a receptor and the high structural requirements for cannabinoid activity indicate the possible presence of a specific endogenous cannabinoid ligand.

SUMMARY OF THE INVENTION

Endogenous ligands for the cannabinoid receptor have not yet been identified. Arachidonylethanolamide, a new arachidonic acid derivative - named anandamide, was isolated from porcine brain. Its structure was determined by mass spectrometry and nuclear magnetic resonance spectroscopy and was confirmed by synthesis. It inhibits the specific binding of a labelled cannabinoid probe to synaptosomal membranes in a manner typical of competitive ligands, and produces a concentration-dependent inhibition of the electrically-evoked twitch response of the mouse vas deferens, a characteristic effect of psychotropic cannabinoids. Similar compounds were synthesized and their pharmacological properties were investigated.

$\Delta^9$ -Tetrahydrocannabinol ( $\Delta^9$ -THC), the psychoactive constituent of Cannabis binds to a specific G-protein coupled receptor in the brain. Although the cannabinoid receptor in the rat and in the human has been cloned, its physiological function is unknown. The well established behavioral effects of THC and the abundance and anatomical localization of the receptor in the brain suggest a role for the receptor in the control of movement, memory, emotions and pain modulation, amongst other activities.

DESCRIPTION OF PREFERRED EMBODIMENTS

The existence of a receptor and the high structural requirements for cannabinoid activity indicate the possible presence of a specific endogenous cannabinoid ligand.

To screen for endogenous cannabinoid compounds there were tested brain-derived fractions using a centrifugation radioligand binding assay. Tritiated HU-243 (11-hydroxyhexahydrocannabinol-3-dimethylheptyl homolog) with a  $K_D$  of 45 pM in rat synaptosomal membranes, was used as the probe. Organic soluble extracts of porcine brain were first chromatographed according to standard protocols for the separation of lipids. Pig brains were homogenized in chloroform and/or methanol and centrifuged at 13,000 x g. The organic soluble extract was fractionated over silica (Kieselgel 60, 70-230 mesh), following elution schemes used to separate the major classes of lipids [C.C.Sweeley, Methods Enzymol. 14, 254 (1969); J.C. Dittmer and M.A. Wells, ibid. p. 482]. Many of the initial fractions isolated from brain inhibited the binding of [ $^3$ H]HU-243 to the cannabinoid receptor. Particular attention was paid to the binding of [ $^3$ H]HU-243 to the siliconized polypropylene microfuge tubes in which the assay was conducted. Normally, about 15-20% of the added [ $^3$ H]HU-243 adheres to the microfuge tube, with the amount increasing slightly when unlabelled cannabinoid drugs

SUBSTITUTE SHEET (RULE 26)

displace the radioligand from the receptor. When monitoring the three-way equilibrium of [ $^3\text{H}$ ]HU-243 among the synaptosomal receptors, the solution and the microfuge tube, we observed that all the crude brain fractions which inhibited the binding of [ $^3\text{H}$ ]HU-243 to the receptors also inhibited the binding of the radioligand to the microfuge tube. Several promising fractions were purified using both low and medium pressure column chromatography as well as thin layer chromatography (tlc). A combination of normal phase and reverse phase systems was employed. Tlc: Rf 0.65 on an analytical RP-18 plate (Merck), eluted with methanol-dichloromethane, 4:1: developed twice-first solvent front: 3.1 cm, second solvent front: 7.4 cm. [W.A. Devane, L. Hanus, Mechoulam, Proceed. 5th Nordic Neuroscience Meet., Publ. Univ. Kuopio Med. p. 198 (1991)]. Anandamide elutes from a silica column (Kieselgel 60, 40-63  $\mu\text{m}$ , Merck) with methanol: chloroform (2:98). It elutes from a reverse-phase column (RP-C, 40-63  $\mu\text{m}$ , Sigma) with methanol: water (88:12).

A compound was recovered (0.6 mg from 4.5 kg of brain), named anandamide, which shows one spot on TLC and elutes mainly as one peak on gas chromatography (GC) using a mass spectrometer as a detector. Anandamide inhibits the specific binding of [ $^3\text{H}$ ]HU-243 to synaptosomal

membranes in a manner typical of competitive ligands with a  $K_D$  of  $52 \pm 1.8$  nM ( $n=3$ ) (Fig. 1). In this system the  $K_D$  of  $\Delta^9$ -THC was  $46 \pm 3$  nM.

Results from previous experiments, in which we compared the inhibitory effects of the 1,1-dimethylheptyl homologs of (+)- and (-)-11-hydroxy- $\Delta^8$ -tetrahydrocannabinol on the electrically-evoked twitch response of the mouse vas deferens, indicate that this preparation is suitable as a model for investigating the mode(s) of action of psychotropic cannabinoids. R.G. Pertwee, L.A. Stevenson, D.B. Elrick, R. Mechoulam, A.D. Corbett, Brit. J. Pharmacol. 105, 980 (1992). Anandamide produced a concentration-dependent inhibition of the twitch response (Fig. 2). The inhibition was not reversed by naloxone (300 nM). The levels of inhibition are comparable to those of binding to the receptor.

The structure of anandamide was established by mass spectrometry (MS) and nuclear magnetic resonance (NMR) spectroscopy. Additional data were obtained from the GC-MS and CID measurements of the trimethylsilyl (TMS) derivative of the material. The results suggest that anandamide is an ethanolamide of a tetraenic  $C_{20}$  fatty acid.

Support for the above structure was found in the behavior of anandamide under GC-MS conditions.

Thermal dehydration gives rise to  $m/z$  329  $M^+$  ion upon electron ionization (EI) and to  $m/z$  330  $MH^+$  under CI. Both self-CI  $m/z$  330  $MH^+$  and  $m/z$  329  $M^+$  are formed under EI conditions in an ion trap instrument (Fig. 3). The fragmentation pattern of the dehydration products was similar in the low mass range of the EI mass spectra of both anandamide and palmitylethanolamide:  $m/z$  85 (McLafferty rearrangement ion) and  $m/z$  98 (product of a  $\delta$ -cleavage) (Fig. 4). The EI mass spectrum of dehydrated palmitylethanolamide exhibits an  $m/z$  112 ion corresponding to a  $\delta$ -cleavage fragment. The absence of this ion in the EI mass spectrum obtained in the GC-MS analysis of anandamide suggests the presence of the first double bond in the tetraenic acid at position 5 (as in arachidonylethanolamide which would not be expected to yield a  $\delta$ -cleavage product) (Fig. 4).

$^1H$  NMR spectra were recorded. The peaks attributed to double bond protons ( $\delta$ , 5.30-5.45, mult) were coupled with those of protons which have the chemical shifts of doubly allylic ones ( $\delta$ , 2.75-2.90, mult). Such doubly allylic protons are typical for numerous naturally occurring all-cis, nonconjugated, polyunsaturated fatty acids such as linoleic and arachidonic acids. Three pairs of protons were observed between  $\delta$  2.01-2.27, which we attributed to



2 allylic methylene groups and 1 methylene group  $\alpha$  to a carbonyl moiety. Only one methyl group was observed (0.88, t). The peaks observed for 2 protons at 3.42 (N-CH<sub>2</sub>, t), 2 protons at 3.72 (O-CH<sub>2</sub>, t) and 2 protons at 2.20 (COCH<sub>2</sub>, t) are similar in chemical shifts and spin coupling patterns to peaks observed in the NMR spectrum of synthetic palmitylethanolamide. The peaks for N-CH<sub>2</sub> and O-CH<sub>2</sub> were coupled.

A juxtaposition of the above analytical data led us to conclude that the structure of anandamide is that of arachidonyl ethanolamide [5,8,11,14-icosatetraenamide, (N,-2-hydroxyethyl)-(all-Z)] a novel chemical entity. This conclusion was confirmed by synthesis. Arachidonyl chloride, prepared from arachidonic acid and oxalyl chloride (21), in methylene chloride, was added at 0 degrees C, under a nitrogen atmosphere to ethanolamine (in a ten fold molar excess) in methylene chloride. After 15 min the reaction was washed with water, dried, and the product (ca 90% yield) was purified by silica column chromatography (eluted with 2% methanol in chloroform) to give arachidonyl ethanolamide, an oil, in 97% purity (by GC-MS). Synthetic arachidonyl ethanolamide was identical with the product obtained on tlc (10), NMR (300 MHz) and GC-MS (retention time and fragmentation

pattern) (Fig. 3). Synthetic anandamide binds to the cannabinoid receptor  $K_i = 39 \pm 5.0$  nM (n=3).

The novel purified compound anandamide seems to be present as brain constituent. It is possible that this compound  
5 is present as a complex with another compound or in any other form, but according to the present invention it has been established that the compound defined herein as ananamide is characterized by the properties set out herein. Similar compounds, defined herein, are  
10 characterized by essentially equivalent properties and are part of the present invention.

This invention also relates to tagged form of such compounds, which can be used in a variety of research projects and in assays.

15 The invention further relates to pharmaceutical compositions containing an effective quantity of one of the compounds defined herein.

Following the synthetic method described for arachidonylethanolamide, the ethanolamides of the following  
20 unsaturated fatty acids were prepared.

	<u>Systematic Name</u>	<u>Trivial Name</u>	<u>Shorthand designation</u>
	9,12-octadecadienoic*	linoleic	18:2 (n-6)
	6,9,12-octadecatrienoic	$\delta$ -linolenic	18:3 (n-6)
5	8,11,14-eicosatrienoic	homo- $\delta$ -linolenic	20:3 (n-6)
	4,7,10,13,16-docosapentaenoic	-	20:5 (n-6)
	9,12,15-octadecatrienoic	$\delta$ -linolenic	18:3 (n-3)
10	5,8,11,14,17-eicosapentaenoic	-	20:5 (n-3)
	4,7,10,13,16,19-docosahexaenoic	-	22:6 (n-3)
	5,8,11-eicosatrienoic	-	20:3 (n-9)

\* The double-bond configuration in each instance is cis.

15 These ethanol amide derivatives have antiinflammatory analgetic, antiglaucoma and antiemetic activity (see Biological Results). In order to obtain  $^{14}\text{C}$  labeled compounds of importance in biological studies the above described synthetic procedure was repeated with  $^{14}\text{C}$ -labeled

20 arachidonic acid to give  $^{14}\text{C}$ -arachidonyl ethanolamide, 50 mCi/mmol. With  $^3\text{H}$ -arachidonic acid we obtained  $^3\text{H}$ -arachidonylethanol amide, 150 Ci/mmol. When we used  $^{14}\text{C}$ -labeled ethanolamide we obtained  $^{14}\text{C}$ -arachidonylethanolamide, 50 mCi/mmol.

## 25 Biological Results

Analgesia. The novel compounds were tested in the standard hot plate test.

Representative results are presented in Table I.

The compounds were dissolved in a detergent (Emulphor):  
ethanol: saline (5:5:90) and administered by intravenous  
injection in the tail vein with an injection volume of  
5 0.1 ml/10g of body weight.

#### Antiemetic activity

The compounds also tested in pigeons against emesis caused  
by an antieoplastic drug (cisplatin) according to the  
method described in Feigenbaum et al., Eur. J. Pharmacol.  
10 169, 159-165 (1989). Representative results are presented  
in Table I.

The compounds were dissolved as described above for the  
tests on analgesia, and administered subcutaneously.  
The injection volume was 1.0 ml/kg body weight.

#### 15 Antiglaucoma activity

The compounds were also tested for antiglaucoma activity  
in rabbits with stable glaucoma induced by  $\delta$ -chymotrypsin  
injection into the eye as described in detail in R.  
Mechoulam et al., On the therapeutic possibilities of some  
20 cannabinoids in "The Therapeutic Potential of Marihuana"  
(eds S. Cohen. R.C. Stillman) Plenum Press, New York,  
1975. pp. 35-48). The compounds were compared in activity  
to that of pilocarpine (a standard antiglaucoma drug)  
which at 0.01% aq. solution reduces the intraocular  
25 pressure (I.O.P.) and delays recovery to half time of the

original I.O.P. about 30 hours.

The fatty acid ethanol amides administered to the eye in 0.1% solution were less active than pilocarpine and delayed recovery to half time of the original I.O.P between 2 and 5 10 hours.

The compounds were dissolved as described for analgesia and further diluted with saline to obtain the required concentration.

#### Antiinflammatory activity

- 10 The compounds were tested by a method reported by W. Calhoun et al., Agents and Actions, 21, 306-a309 (1987). Water was substituted for mercury as the displacement medium. PAF (1.0 µg) or arachidonic acid (1.0mg) dissolved in 50 µl of 5% ethanol in saline was injected 15 s.c. into the plantar surface of the right hind paw of female mice (20-25g). The mice were under ether anesthesia during this procedure. The volume of the right foot was measured to the level of the lateral malleolus by water displacement before treatment and 15 min after 20 PAF injection or 30 min after arachidonic injection. The change in paw volume was calculated for each mouse. The results for two of the tested ethanolamides are presented in Table II.

Dosage

The effective doses for humans are between 1-100 mg total daily dose, by injection or by oral administration.

Table I

5

Analgesia and Vomiting Reduction

<u>Compound</u>	<u>Analgesia</u> ED <sub>50</sub> (mg/kg)-	<u>Reduction of</u> <u>vomiting(50%)=</u> (mg/kg)
Arachidonylethanolamide	4.2	2.5
10 5,8,11,14,17-eicosapentaenyl ethanolamide	5.2	3.6
4,7,10,13,16,19-docosa- hexaenylethanolamide	6.1	4.8
15 5,8,11-eicosatrienyl- ethanolamide	7.6	6.2

- in mice. For details of administration see Text  
= in pigeons. For details of administration see Text.

Table II

Inhibition of Arachidonic Acid-Induced Paw Edema-

b			
5	Dose (mg/kg)	Arachidonyl-	4,7,10,13,16,19
		ethanolamide	Docosahexaenyl ethanolamide
	0.010	56.2	-
	0.025	58.4	-
10	0.050	74.2	52.4
	0.100	98.0	66.8
	0.250	100.0	72.0

= Values shown are percent inhibition of paw edema when compared to vehicle treated controls. 95% significance by ANOVA. N=5 mice/group.

b Control mice were given peanut oil (50ul) orally. Paw volume increase =  $38 \pm 4 \mu\text{l}$ .

BRIEF DESCRIPTION OF THE FIGURES

The invention is illustrated with reference to the enclosed Figures, wherein:

- Fig. 1 illustrates the competitive inhibition of [ $^3\text{H}$ ] HU-243 binding by natural anandamide;
- Fig. 2 illustrates inhibition of vas deferens twitch response by natural anandamide;
- Fig. 3 is a GC-MS spectrum of anandamide;
- Fig. 4 illustrates the structure of some compounds described;
- Fig. 5 EI GC-MS spectra of dehydrated homo- $\delta$ -linolenylethanolamine (5);
- Fig. 6 EI GC-MS spectra of dehydrated docosatetraenylethanolamine, Compound (6);
- Fig. 7 EI GC-MS spectra of homo- $\delta$ -linolenylethanolamine (5) and of docosatetraenylethanolamine, (top and bottom);
- Fig. 8 illustrates competitive binding of [ $^3\text{H}$ ]HY-243 by compound (5), squares, and by compound (6), triangles;
- Fig. 9 is a Formula sheet of some compounds of the invention.



Legends

Fig. 1. Competitive inhibition of [ $^3$ H]HU-243 binding by natural anandamide. Synaptosomal membranes were prepared from rat whole brain-minus brainstem (Sprague-Dawley males, 430-470g). [ $^3$ H]HU-243 (45-55 pM) was incubated with synaptosomal membranes (protein content 3-4 ug) for 90 min at 30 C with either the indicated concentrations of anandamide or the vehicle alone (fatty acid-free bovine serum albumin, final concentration-0.5 mg/ml). Bound and free radioligand were separated by centrifugation (see ref. 8 for full details of the assay). The data were normalized to 100% of specific binding, which was determined with 50 nM unlabeled HU-243. The amount of specific binding was 77-82% of the total radioactivity bound to the membranes. Data points ( $\square \Delta$ ) represent the average of triplicate determinations from two independent experiments.  $K$  value (mean + SE, n=3) was determined using the Ligand program;  $K = 52 \pm 1.8$  nM.

Fig. 2. Inhibition of vas deferens twitch response by natural anandamide. Vasa deferentia obtained from MFI mice were mounted in 4 ml siliconized organ baths under an initial tension of 0.5 g. The baths contained

Mg<sup>++</sup> -Free Krebs solution which was kept at 37 C and bubbled with 95% O<sub>2</sub> and 5% CO<sub>2</sub>. Tissues were stimulated supramaximally with 0.5 s trains of 3 pulses (train frequency 0.1 Hz: pulse duration 0.5 ms). Isometric  
5 contractions were recorded. Natural anandamide was dispersed in Tween 80 and saline (see ref. 12) and added in volumes of 40 ul. Tween 80 did not inhibit the twitch response at the maximum bath concentration used (0.63 ug/ml; n=6).

10 Fig. 3. GC-MS spectrum of anandamide on an ion trap instrument. Anandamide undergoes thermal dehydration under these conditions (see text) hence the above spectrum in actuality is that of the M<sup>+</sup> ion of the corresponding 2-oxazoline.

15 Fig. 4. Structures of anandamide and of palmitylethanolamide and of the dihydro- and tetrahydrooxazole ion fragments formed from these fatty acid ethanolamides on thermal dehydration under GC-MS  
20 conditions (see text). The m/z 112 ion is formed only from palmitylethanolamide.

## COMPOUNDS ISOLATED FROM THE BRAIN AND SYNTHESIZED.

<u>NAME OF COMPOUND</u>		<u>K<sub>i</sub></u>		
	cis-7,10,13,16-docosatetraenoy-			
	lethanolamide	34.4	± 3.2	nM
5	anandamide	39.0	± 5.0	nM
	homo- $\gamma$ -linolenoylethanolamide	53.4	± 5.5	nM

## SYNTHETIC COMPOUNDS, PREPARED AND TESTED.

<u>NAME OF COMPOUND</u>		<u>K<sub>i</sub></u>		
	N-propyl-5,8,11,14-eicosa-			
10	tetraenoylamide	11.7	± 2.6	nM
	N-ethyl-5,8,11,14-eicosa-			
	tetraenoylamide	34.0	± 2.7	nM
	N-methyl-5,8,11,14-eicosa-			
	tetraenoylamide	60.0	± 7.4	nM
15	arachidonoyl- $\beta$ -dimethy-			
	lethanolamide	161.8	± 34.1	nM

## Analysis:

## Compound 5:

5 Calcd for  $C_{22}H_{39}NO_2$  (m.w. 349) C, 75.59  
H, 11.25  
N, 4.01

Found C, 75.85  
H, 11.01  
N, 3.88

## Compound 6:

10 Calcd for  $C_{24}H_{41}NO_2$  (m.w. 375): C, 76.75  
H, 11.00  
N, 3.73


15 Found C, 76.92  
H, 11.34  
N, 3.60

Appendix II Active Unsaturated Acylethanolamides Isolated and/or Prepared.

Name of the compound  $K_1$  (binding to receptor)


cis-5,8,11,14-eicosatetraenyl ethanolamide  
(anandamide)  
from 20:4 (n-6) acid  
synthetic and from brain

$39.0 \pm 5$  nM




cis-7,10,13,16-docosatetraenylethanolamide  
(from 22:4 (n-6) acid)  
synthetic and from brain

$34.4 \pm 3.2$  nM




homo- $\gamma$ -linolenylethanolamide  
(cis-8,11,14-eicosatrienylethanolamide)  
(from cis-8,11,14-eicosatrienoic acid) 20:3 (n-6)  
synthetic and from brain

$53.4 \pm 5.5$  nM




cis-4,7,10,13,16,19-docosahexaenylethanolamide  
from 22:6 (n-3) acid

328.9 nM




cis-11,14-eicosadienylethanolamide  
from 20:2 (n-6) acid

1.5  $\mu$ M



$\gamma$ -linolenylethanolamide  
(cis-6,9,12-octadecatrienylethanolamide)  
(from cis-6,9,12-octadecatrienoic acid) 18:3 (n-6)

$4.6 \pm 0.3$   $\mu$ M




Pharmaceutical compositions, containing as active ingredient an effective quantity of the compounds of the present invention, have a variety of pharmacological effects. Amongst these there may be mentioned anti-inflammatory, anti-asthmatic, analgetic, antiglaucoma, anti-migraine and anti-spasticity effects. They are also mood-stimulating and ameliorate the symptoms of multiple sclerosis. The unit dosage form varies according to the compound and medical use, and is generally in the range between about 1 mg to about 100 mg. The preferred range is between about 5 to about 25 mg per unit dosage form.

## Claims :

21

1. Essentially pure compounds of general formula



- wherein R is the alkenyl moiety of a polyunsaturated fatty acid of 16 to 28 carbon atoms, with 2 to 6 double bonds, preferably all in the cis- configuration, with the first double bond at the C-3, C-6, or C-9 position, counting from the non-carboxyl part of the molecule, where R'' is -H, lower-alkyl, -OH or  $-(\text{CH}_2)_n\text{-OH}$ , where n is a small integer, and
- when R'' is hydrogen, R' is lower alkyl, or  $-(\text{CH}_2)_m\text{OH}$ , where m is a small integer,
- when R'' is lower-alkyl, R' is  $-(\text{CH}_2)_p\text{-OH}$ , where p is a small integer, when R'' is -OH, R' is  $-(\text{CH}_2)_q\text{-OH}$  where q is a small integer, or both R' and R'' are each  $-(\text{CH}_2)_n\text{-OH}$ , where n is a small integer, and acid addition salts and complexes of these.

2. A compound according to claim 1, where the alkenyl moiety is a octadecadienoic, actadecatrienoic, eicosapentaenoic, docosaheptaenoic eicosatrienoic or eicosatetraenoic moiety.

3. A compound according to claim 1, selected from arachidonylethanolamide, cis-7,10,13,16-docosatetraenylethanolamide, homo- $\delta$ -linolenylethanolamide, N-propyl-5,8,11,14-eicosatetraenoylamide, N-ethyl-5,8-11, 14-eicosatetraenoylamide, N-methyl-5,8,11,14-eicosatetraenoylamide arachidonoyl- $\beta$ -dimethylethanolamide.

4. An antiinflammatory, anti-asthmatic, analgetic, anti-emetic, antiglaucoma, anti-migraine, anti-spasticity, mood stimulating and symptoms of multiple sclerosis ameliorating pharmaceutical composition, which contains as  
5 active ingredient an effective quantity of a compound defined in claim 1, 2 or 3.
5. A composition according to claim 4, where the active ingredient is a compound defined in claim 3.
6. A composition according to claim 4 or 5, where the  
10 unit dosage for human administration is from about 1 mg to about 100 mg of the active compound.
7. A compound defined in claim 1, 2 or 3 in radioactive tagged form.
8. A compound according to claim 7, where the tag is  
15  $^3\text{H}$  or  $^{14}\text{C}$ .
9. A composition for binding to the cannabinoid receptor in brain, comprising an effective quantity of a compound defined in any of claims 1, 2 or 3.



1/9




FIG. 1

2/9




FIG. 2

3/9




FIG. 3

4/9







FIG. 4

5/9




FIG. 5

SUBSTITUTE SHEET (RULE 26)




FIG. 6

7/9




FIG. 7

SUBSTITUTE SHEET (RULE 26)

8/9




FIG. 8



9/9




FIG. 9

SUBSTITUTE SHEET (RULE 26)

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US93/11625

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) : C07C 233/00

US CL : 554/66; 514/627,826,886,887,903,

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 554/66; 514/627,826,886,887,903

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, CAS Online

search terms: anandamide, MS/multiple sclerosis, migraine, cannabinoid receptor, glaucoma, anti-inflammatory/asthmatic/emetic

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X — Y	US, A, 4,877,789 (Shroot et al) 31 October 1989, column 1, lines 6-12, 37-47, 62-67; column 2, lines 1-18, 62-67; column 3, lines 1-10; claims 1-4, 6-11, 13.	1-6 — 1-6
X — Y	US, A, 4,619,938 (Takahashi et al) 28 October 1986, column 1, lines 54-68; column 2, lines 1-2, 7-18; claims 1-4.	1-6 — 1-6

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* "A"	Special categories of cited documents: document defining the general state of the art which is not considered to be part of particular relevance	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E"	earlier document published on or after the international filing date	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O"	document referring to an oral disclosure, use, exhibition or other means	"A"	document member of the same patent family
"P"	document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

02 MARCH 1994

Date of mailing of the international search report

15 MAR 1994

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. 703-305-3230

Authorized officer

DEBORAH D. CARR

Telephone No. 703-308-1235